



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Improved performance in master runners competing in the European Championships between 1978 and 2014

Schneider, Anaïs L ; Nikolaidis, Pantelis Theodoros ; Knechtle, Beat

Abstract: The performance trends in elite runners have been well investigated, but we have no knowledge about performance trends and the difference between the sexes in elderly runners competing at a high level in varying distances. The purpose of this study was to investigate the performance of these age groups. Data from seventeen European Championships held between 1978 and 2014 were analyzed for various race distances (i.e. 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, 10,000 m and marathon). Running speed for the top 8 female and male finalists for each age group (35-99 years, split into 5 year gaps) and each race distance were included. A two-way analysis of variance compared the effects of sex, race distance, age group and calendar year on speed. Subsequent comparisons between race distances, age groups or calendar years were carried out using a post-hoc Bonferroni test. Our analysis shows that men were faster than women in all distances and the difference between the sexes was greater in the shorter distances. Speed was higher for shorter distances than for longer distances. Younger participants were faster than older ones, and the effect of age group was the largest for the 200 m. There was a minor effect of calendar year on speed in the 100 m, 200 m, 1500 m, 10,000 m and marathon, and a minor calendar year×sex interaction on running speed was shown for the 200 m. For athletes and coaches, the current study demonstrates that both male and female athletes improved their running performance over time and that the sex gap may have reached its limit.

DOI: <https://doi.org/10.1519/JSC.0000000000002548>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-152639>

Journal Article

Accepted Version

Originally published at:

Schneider, Anaïs L; Nikolaidis, Pantelis Theodoros; Knechtle, Beat (2019). Improved performance in master runners competing in the European Championships between 1978 and 2014. *Journal of Strength and Conditioning Research*, 33(9):2559-2569.

DOI: <https://doi.org/10.1519/JSC.0000000000002548>

Improved Performance in Master Runners Competing in the European Championships between 1978 and 2014

Performance trends in master runners grouped by age

Anaïs L. Schneider ¹, Pantelis T. Nikolaidis ², Beat Knechtle ^{1,3}

¹Institute of Primary Care, University of Zurich, Zurich, Switzerland

²Exercise Physiology Laboratory, Nikaia, Greece

³Medbase St. Gallen Am Vadianplatz, St. Gallen, Switzerland

Corresponding author:

Prof. Dr. med. Beat Knechtle

Facharzt FMH für Allgemeinmedizin

Medbase St. Gallen Am Vadianplatz

Vadianstrasse 26

9001 St. Gallen

Switzerland

Telefon +41 (0) 71 226 93 00

Telefax +41 (0) 71 226 93 01

E-Mail beat.knechtle@hispeed.ch

1 Abstract

2 The performance trends in elite runners have been well investigated, but we have no
3 knowledge about performance trends and the difference between the sexes in elderly
4 runners competing at a high level in varying distances. The purpose of this study was
5 to investigate the performance of these age groups. Data from seventeen European
6 Championships held between 1978 and 2014 were analyzed for various race distances
7 (i.e. 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, 10,000 m and marathon).
8 Running speed for the top 8 female and male finalists for each age group (35-99
9 years, split into 5 year gaps) and each race distance were included. A two-way
10 analysis of variance compared the effects of sex, race distance, age group and
11 calendar year on speed. Subsequent comparisons between race distances, age groups
12 or calendar years were carried out using a post-hoc Bonferroni test. Our analysis
13 shows that men were faster than women in all distances and the difference between
14 the sexes was greater in the shorter distances. Speed was higher for shorter distances
15 than for longer distances. Younger participants were faster than older ones, and the
16 effect of age group was the largest for the 200 m. There was a minor effect of
17 calendar year on speed in the 100 m, 200 m, 1500 m, 10,000 m and marathon, and a
18 minor calendar year \times sex interaction on running speed was shown for the 200 m. For
19 athletes and coaches, the current study demonstrates that both male and female
20 athletes improved their running performance over time and that the sex gap may have
21 reached its limit.

22

23 **Key words:** speed; runner; sex difference; age group athlete

24 INTRODUCTION

25 Track and road running is an increasingly popular sports discipline, where participants
26 aim to cover a given distance – usually from 100 m to marathon – as fast as possible
27 (62). The last few years have seen an increase in participation, especially in marathon
28 road running (21, 37). Despite the growing participation in running events of varying
29 distances, little is known about the performance trends of elderly female and male
30 elite athletes in track running (49).

31 Running speed has improved considerably for both women and men during the last
32 150 years (62). World records in all running events have been continually beaten over
33 the past century (41). Several factors might contribute to the dramatic increase in
34 running performance. In the late 1970s sports physiology began to be applied to sports
35 training strategies. Different training methods were incorporated, including circuit
36 training, interval training and sprints. A better understanding of various training
37 components probably played a major role in the increase of performance in the last
38 decades (59). Incorporation of an interval training session has been proven to increase
39 running speed on longer runs and lengthen the time to exhaustion in trained athletes
40 (32). During the last century, studies started to include research about resistance
41 training. The first studies about resistance training started in the 1980s. Since then it
42 has been shown that resistance training has an impact on neural mechanisms,
43 metabolic adaptations, the cardiovascular system, the endocrine system, connective
44 tissue and the immune system (33). Changes in training strategies such as an
45 increased training volume helped achieve this progress (52). A greater interest in sport
46 psychology might have helped understanding an athlete's psychological needs in
47 order to perform better (34). Keegan et al. studied the impact of coaches, parents and
48 peers on an athlete's motivation and showed that the behaviour of the people in an

49 athlete's environment has a direct, but complex impact on his or her motivation (25).
50 Coaching strategies have changed over time and provide athletes with better
51 techniques for coping with anxiety, which can negatively affect athletic performance
52 (44).

53 Progress in the medical field might also play a role as it provides more concise
54 medical advice in terms of prevention and therapy of sport injuries (59). For example,
55 research about overtraining syndrome in athletes helps understand the aetiology of
56 this condition and how to avoid and treat it (6). Radiographic imaging, arthroscopy
57 and new surgical methods help provide a better, more complete medical support and
58 play a major role in the increase of performance (59). Another contributing factor
59 might be found in the better understanding of the nutritional demands of an athlete's
60 body (59) and of the hydration needed during endurance performance (26). Advances
61 in health technology including the development of functional magnetic resonance
62 imaging have helped understand brain activity during sports (19).

63 In the last decades, considerable advances have also been made in the field of sports
64 equipment. We now have access to improved design and quality of shoes, clothing
65 and training equipment. These improvements also helped athletes achieve an increase
66 in sports performance (59). Unfortunately, doping probably also plays a major role in
67 the increase of athletic performance in the last few decades (41). All of these factors
68 and more could play a role in the overall increase of athletic performance.

69 Men run faster than women. There are several reasons for this difference in endurance
70 performance between the sexes, including higher VO_2max , lower body fat percentage
71 (55), larger hearts, higher haemoglobin levels and greater muscle mass in men (4).
72 Endocrine differences are not to be neglected. Androgens levels are higher in men,

and contribute to their higher muscle mass (51), which has an impact on neuromuscular performance (12). Although both men and women have improved their running performance, the relative improvement in running performance observed over the last century was greater in female than in male athletes (41). Some findings even suggested that women would outrun men at some point in the future (64, 65).

The hypothesis of the narrowing difference between the sexes in running performance has been analysed by other authors (8, 21, 54, 68). For instance, Zingg et al. (69) showed that female athletes over 35 years reduced the difference in mountain and city marathons within a decade. In 1922, the difference was about 30% and decreased over time, reaching a plateau at 10.7% in 1984, according to Thibault et al. (58). Other authors state a similar percentage (15, 21). Interestingly, one study shows that the difference between the sexes is smaller in ultra-marathon running, ranging from 0.2-10% (31). The discussion of whether women will catch up to men or whether the difference might be fixed is still ongoing.

Elderly athletes represent a highly active and healthy population with a good quality of life (67). Thus, studies of performance of this athletic elderly population might help to understand the decline in general health with increasing age (67). Aging is generally associated with a decline in both health (40, 67) and endurance performance (56). Over the last few decades, however, greater numbers of elderly people have participated in endurance races (23, 42, 56). In running, elderly athletes have significantly improved their performance between 1975 and 2013 (2).

Regular endurance training, including running, has many health benefits. Several studies demonstrated the benefits of physical activity later in life such as lower overall mortality (18, 63) Lee et al. analysed the mortality risk of over 13,000 runners and

found out that runners have a lower mortality risk than non-runners (36). It has been shown that regular exercise reduces the incidence of cardiovascular and respiratory diseases (46). Couppé et al. found that elderly men who had been practicing endurance running for a long time had lower triglyceride and LDL-cholesterol than untrained elderly men. The same group analysed accumulation of advanced glycation end products in the connective tissue and showed that elderly elite athletes have a lower AGE accumulation and higher MRI signal intensity of the patellar tendon, suggesting that endurance training could reduce the age-related deterioration of soft tissues (11). Regular physical exercise has also been shown to improve cognitive function (57).

Based on existing reports describing other disciplines like swimming (27) where improvement over time was shown for elderly swimmers in all age groups and distances, we might assume similar trends in running; however, we have no knowledge about changes in performance trends in age group track runners competing at an elite level in various distances. Such knowledge would be of great value from a theoretical and practical perspective. Sports scientists, especially exercise physiologists, focusing on sex differences would benefit from such knowledge, because they would improve their understanding of the evolution of the differences between males and females in sport performance. On the other hand, coaches and fitness trainers working with age group runners might use such information on sex differences to optimize the training of their athletes.

The possibility of a sub-2-hour marathon has been discussed vividly lately. Though runners are getting faster over time, Tucker et al. do not believe in an imminent sub 2-hour marathon because the current average difference between the sexes is about 11.2% and the achievement of a sub 2-hour marathon would represent a 12.9%

difference. Another point they bring forward is that the 2.4% required improvement in men's performance is unlikely to happen quickly, but rather will take generations to achieve (60). Hoogkamer et al. do believe a sub 2-hour marathon is achievable and drew a draft of how a lighter shoe, a downhill route and tailwind might someday reduce the metabolic cost enough in order to achieve the average velocity of 5.86 m/s required for a sub 2-hour marathon (20).

The first aim of this study was to analyse the performance trends in elite master runners from 1978 until 2014 for 5-year age group intervals from 35 to 99 years for different race distances (i.e. 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, 10,000 m and marathon). The second aim was to analyse the difference between men's and women's running performance. We hypothesized that between 1978 and 2014 both men and women improved their running speed and that the difference between the sexes decreased.

METHODS

Experimental Approach to the Problem

In order to investigate the changes of performance (i.e. running speed) of master athletes across the years along with the difference in performance between men and women, we analysed the running speed of men and women for different distances. Data from 17 European Championships held between 1978 and 2014 were analyzed for different race distances (i.e. 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, 10,000 m and marathon). Only outdoor track races and marathons held on roads were included due to the different events (e.g. 100 m versus 60 m) and conditions (e.g. 400 m in one lap versus 400 m in two laps) in outdoor and indoor races, respectively. The

youngest athletes in the data set were 35 years old. Reaburn and Dascombe (49) defined master athletes as participants of 35 years or older. We therefore considered athletes over 35 years of age to be master athletes and did not include any younger athletes in our study.

Subjects

The European Veterans Athletic Association (EVAA) was founded in 1978 in Italy with the aim to organize European Championships for master athletes and to keep a complete register of the results (<http://european-masters-athletics.org/about-us/history.html>). We aimed to investigate the results of the European Championships held between 1978 and 2014. The section www.evaa.ch/results.html records the publicly available race results for the years 2000-2014. The result charts for races held between 1978 and 2000 were not available on the website, thus, Ove Edlund, Sweden, provided us with the printed versions for this time period. Both the online and the printed race records included results for the distances 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, 10,000 m and the marathon. Each athlete was recorded for distance, race time, nationality and age group. Not every Championships result chart included the competitor's exact age, however, every Championships used the same age groups where athletes were divided into 5-year age group intervals as follows: 35-39, 40-44, 45-49, 50-54, 55-59, 60-64, 65-69, 70-74, 75-79, 80-84, 85-89, 90-94, and 95-99 years. To our knowledge no subject was excluded from the EVAA data set because of the use of doping. This study was approved by the Institutional Review Board of Kanton St. Gallen, Switzerland, with a waiver of the requirement for informed consent of the participants as the study involved the analysis of publicly available data (01/06/2010).

Procedures

We analysed the running performance for athletes between 35 and 99 in 5-year age intervals. The race distances included were 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, 10,000 m and marathon. Finally, 14,685 race times (6,019 female and 8,666 male times) were included into the data set. The data set is visible in Table 1. For better comparison, we restricted our analysis to the top eight male and female finishers, as there were different numbers of finishers in each category. Finishers without a recorded race time or who were disqualified for any reason were excluded from our analysis. Running speed was used to compare the performance. As the race times and the distances but not the running speed were listed in the result charts, running speed was calculated in km/h using the equation $\text{running speed (km/h)} = \text{race distance (m)} / \text{race time (sec)} \times 3.6$.

Statistical Analyses

The statistical software IBM SPSS v.23.0 (SPSS, Chicago, USA) performed all statistical analyses. Mean values and standard deviation (s) were calculated for all variables. A two-way analysis of variance (ANOVA) compared effects of sex, race distance, age group and calendar year on running speed. Subsequent comparisons among race distances, age groups and calendar years were carried out using post-hoc Bonferroni test. The magnitude of these differences was examined using effect size eta squared (η^2) and evaluated as: minor ($0.010 < \eta^2 \leq 0.059$), moderate ($0.059 < \eta^2 \leq 0.138$) and major ($\eta^2 > 0.138$) (9). We analysed the relationship between speed and race duration using a logarithmic regression model. In addition, we examined the differences in the running speed of men and women using the formula

100×(men's running speed–women's running speed)/women's running speed. We also compared variations in running speed by participants' sex, age group, race distance and calendar year by a mixed-effects regression model. In this model, participants were assigned as random variables, whereas sex, age group, race distance and calendar year were assigned as fixed variables. We examined interaction effects among these fixed variables. Akaike information criterion (AIC) was used to select the final model. These analyses were performed for each race distance separately. A regression analysis of cubic degree was performed between running speed and calendar year, and the coefficient of determination (R^2) was calculated. Statistical significance was set at $\alpha = 0.05$.

RESULTS

Performance by sex and race distance

According to the two-way ANOVA, a moderate effect of sex on running speed was observed ($p < 0.001$, $\eta^2 = 0.115$), where men were faster than women, as shown in Figure 1. Also, a major effect of race distance on running speed was shown ($p < 0.001$, $\eta^2 = 0.649$), where short distances were faster than longer ones. In addition, a sex×race distance interaction on running speed was noticed ($p < 0.001$, $\eta^2 = 0.006$) for all distances with the sex difference being greater in the shorter distances. These findings were in agreement with the mixed-effects regression analysis and are shown in Table 2. Figure 2 presents the relationship between running speed and race duration.

Performance by sex and age group

A major effect of sex on running speed ($p < 0.001$), where men were faster than women, was observed for all distances with η^2 ranging from 0.197 (1000 m) to 0.472 (200 m). Figure 3 represents the difference between the sexes for each distance and each age group. In addition, a major effect of age group on running speed ($p < 0.001$), where the younger groups were faster than the older, was shown for all distances with η^2 ranging from 0.474 (marathon) to 0.823 (200 m). A minor sex \times age group interaction on running speed ($p < 0.001$), where difference between the sexes was greater in the younger age groups, was noticed for all distances with η^2 ranging from 0.019 (1500 m) to 0.048 (100 m). The findings of the mixed-effects regression analysis were shown in Table 3.

Performance by sex between 1978 and 2014

A minor effect of calendar year on running speed was observed in 100 m, 200 m, 1500 m, 10,000 m and marathon ($p \leq 0.044$, $0.014 \leq \eta^2 \leq 0.050$), but not in 400 m, 800 m and 5000 m ($p \geq 0.160$, $\eta^2 \leq 0.011$) (Figure 4). A small calendar year \times sex interaction on running speed ($p = 0.038$, $\eta^2 = 0.013$) was shown for 200 m, but not for the rest of the distances. The findings of the mixed-effects regression analysis were presented in Table 3. The regression analysis of cubic degree between sex difference and calendar year showed variation by race distance: 100 m ($R^2 = 0.41$), 200 m ($R^2 = 0.34$), 400 m ($R^2 = 0.628$), 800 m ($R^2 = 0.309$), 1500 m ($R^2 = 0.209$), 5000 m ($R^2 = 0.328$), 10,000 m ($R^2 = 0.489$) and marathon ($R^2 = 0.046$). These findings are shown in Table 4 and Figure 5.

241 **DISCUSSION**

242 The main findings of the present study were that (i) men were faster than women in
243 all distances, (ii) the difference between men and women in running speed was greater
244 in the shorter distances, with the greatest difference in 200 m and the smallest in
245 10,000 m, (iii) running speed was faster for shorter distances than for longer
246 distances, (iv) younger participants were faster than older ones, (v) with the effect of
247 age group the largest in 200 m and the smallest for the marathon, (vi) there was a
248 minor effect of calendar year on running speed in 100 m, 200 m, 1500 m, 10,000 m
249 and marathon, and (vii) a minor calendar year \times sex interaction in running speed was
250 shown for 200 m.

251

252 **Men were faster than women for all distances and age groups**

253 The first result was that men were faster than women for all distances and age groups,
254 which was supported by other studies (21, 54). Generally speaking, the difference of
255 10-12 % between male and female running speeds has been consistent for both shorter
256 and longer distances (16).

257 Although one might argue that this result was trivial, recent studies investigating
258 master pool swimmers in freestyle (27), breaststroke (29), backstroke (61) and open-
259 water swimmers (28) showed that men were faster than women for younger age
260 groups, but not for older age groups where women achieved a similar performance to
261 men. The explanation was that in the studies with swimmers at the World
262 Championships, no selection by age group occurred and all swimmers in all age
263 groups were considered. The men-to-women ratio changed across age groups and,

therefore, explains the fact that women were able to achieve a similar performance to men in the older age groups.

Men were faster than women in our study. This may be due to their lower body fat percentage and their higher VO_2max (55) compared to women. However, these hypotheses were limited, as we did not perform any determinations of body fat or VO_2max . Men also seem to be more competitive (13) and more willing to commit to a high intensity training program (16) than women. There might also be a historical component as it is known that women have been participating in official races for a shorter period of time and have therefore less experience in running races (21). We did not collect any historical data ourselves, but it is known that the first modern Olympic Games were introduced in 1896 with only male athletes, and a century later (in 1984), female runners were first included for the marathon distance (59).

There are other differences between male and female runners. Women tend to have a more even pace throughout a race, while men slow down for the second half of the marathon (14). This difference might be due to different choices while racing or to the fact that men are depleting their muscle glycogen more rapidly than women (14).

In the present study we used running as a way to compare male and female performance and the changes over time. Distance running is a suitable discipline to compare male and female performance because it is objective, open to everyone who wants to participate and popular among men and women (13). Sex differences in endurance performance were also found in other athletic disciplines. In swimming races, the sex differences seem more important in elderly than in younger swimmers, though not as important as in marathon running (53).

The difference between the sexes was greater in shorter race distances

A second important discovery was that the difference in male and female running speed was greater for shorter distances and smaller for longer distances. The effect of sex was the highest for the 200 m distance and the smallest for the 10,000 m distance. Sprint performance depends on peak power output, which increases linearly with the amount of lean mass in the lower extremities, and is higher in males (43). Proportionally, women oxidize more fat than men during endurance exercise (7), which might help explain why women are less fatigued on longer distances (8). Other contributing factors to fatigue could be hormonal differences as oestrogens are known to have a protective role in exercise-related muscle damage that accompanies longer aerobic exercise (66).

Running speed was faster for shorter distances

Another discovery was that running speed was faster for shorter distances than for longer ones. This result was expected and has been demonstrated by other authors. For example, Weiss et al. (62) showed that running speed decreased with increasing race distance. Different physiological factors cause a reduction of physical effort including cardiorespiratory capacity, $\text{VO}_{2\text{max}}$ and muscle fatigue (24).

Younger athletes were faster than elderly athletes

Another important result was that younger competitors ran faster than elderly athletes for every distance. This effect of age on running speed was expected and has been demonstrated by other authors (10, 21, 30, 56).

The best running times are often achieved by the age of 35 years with a small and linear decline until about 50-60 years of age, with a more pronounced decline after that (56). A decline in VO_2 which is due to a reduction in cardiac output, an increase in body fat percentage, a reduction in peripheral oxygen extraction and most importantly, a reduction in muscle mass (48), seems to play an important role in the decline of running performance in elderly athletes (47). Younger master athletes have higher muscle mass and a greater number of motor units than older master athletes (17). Muscle atrophy causes a reduction of muscle mass (1, 3). Muscle atrophy in turn could partially be explained by neuronal dysfunction (5). Therefore, the age-related muscle loss contributes to the reduction of $\text{VO}_{2\text{max}}$ (48) which in turn plays an important role in the age-related decrease of running speed (47).

The oldest competitors in our study were 97 years old, but other authors reported great sports performance of centenarians (38), showing that physical activity is possible until late in life. Age-related sarcopenia could also play a role in the performance decline. Sarcopenia is defined by a decrease in number and function of muscle fibres. It is the result of various factors including age-related defect in autophagy resulting in protein accumulation, impaired mitochondrial function due to increased reactive oxygen species, diminished regenerative potential of muscle fibres, degenerative atrophy, vitamin deficiencies and hormonal changes. Muscle mass and strength are also linked to testosterone levels. An increase in age correlates to decreased testosterone levels in both men and women, which is linked to a decrease in muscle mass, particularly in men (45)

The effect of age group was greatest for 200 m and smallest for the marathon

Another finding was that the age-associated decline in performance was the largest for the 200 m distance and the smallest for the marathon. There are divergent results in current literature. Rittweger et al. showed that the age-related decrease in running performance was similar for both sprint and long-distance runners (50). Drey et al. (17) compared the skeletal muscle mass in power- and endurance-trained master athletes competing in the 2012 European Championship held in Zittau, Germany. Power-trained master athletes had a better muscle mass than endurance-trained master athletes (17). During the aging process relatively more type II muscle fibres are lost (39). This could explain why in our study, the age-related decline was more pronounced for the 200 m distance and smallest for the marathon. Both sprinting and long-distance training are beneficial during the aging process (35). The fact that 99-year old athletes were competing in Championships shows that aging might come with a decrease in running performance but it is compatible with an active lifestyle.

Effect of calendar year on speed in 100 m, 200 m, 1500 m, 10,000 m and marathon

Another important discovery was an effect of calendar year on running speed for most distances. It has been suggested by different authors that running performance has been improving over the years (56, 62). World records in athletic sports like running have dramatically increased over the past century. The improvements in race times were directly proportional to race distances, meaning the smallest improvements were achieved on short distances like 100 m and the largest improvements were observed in

longer distances such as marathon running (41). Weiss et al. showed that the improvement observed at the beginning of the century is now stagnating (62).

Small calendar year×sex interaction on speed for 200 m

The last significant finding was that the difference between the sexes has decreased over the years, but only for the 200 m distance. We expected that the difference would be decreasing for every distance; however, results of earlier studies were divergent. Thibault et al. (58) showed that the difference has been stable since 1984 and that the gap will most likely not be closed. Hunter et al. (22) show that the difference diminished between 1980 and 2010. The researchers also claimed that the male-female difference is closely linked to the male-female participation ratio and therefore many studies have a historical and sampling bias (21).

Limitations and strengths

A limitation of these discoveries was that the present study focused on European athletes and outdoor Championships. Thus, caution was needed to generalize these findings to non-European athletes and indoor Championships. Non-Europeans showed superior performance in athletic running events and their performance would be expected to vary differently by calendar year and sex. Moreover, some events differed between outdoor and indoor competitions (e.g. 100 m versus 60m, 5000 m versus 3000m, respectively) or do not exist in indoors competitions (10,000 m and marathon) and those which existed in both competitions (e.g. 400 m) included different numbers of laps (e.g. one versus two) affecting the physiological, technical and tactical demands of a similar distance. Nevertheless, the strength of the study was its novelty

as it is the first one in this topic, adding novel information about the variation of performance of master runners by calendar year and sex.

In conclusion, male runners are generally faster than females. The the gap between male and female endurance performance has been decreasing over the past decades but that it seems to have reached a limit. In general, all athletes are getting faster over time.

PRACTICAL APPLICATIONS

For athletes and coaches, information acquired through this study could have practical implications for practitioners working with master runners; most training programs were developed and applied originally in men, and, thus, knowledge about sex differences would help coaches designing tailored training programs for women, as well. In addition to this practical application, the results about the variation of performance by age group and the variation of the effect of age by race distance would be of interest for researchers studying master athletes as a model of effective aging and for sports scientists, especially exercise physiologists, focusing on sex differences in sport performance.

Acknowledgements

We thank Ove Edlund, Sweden, for providing us the data before the year 2000 and Mrs Knechtle for transcribing this data set. We also thank Patricia Villiger for her help in translation.

404 **References**

- 405 1. Akima H, Kano Y, Enomoto Y, Ishizu M, Okada M, Oishi Y, Katsuta S, and
406 Kuno S. Muscle function in 164 men and women aged 20--84 yr. *Med Sci*
407 *Sports Exerc* 33: 220-226, 2001.
- 408 2. Akkari A, Machin D, and Tanaka H. Greater progression of athletic
409 performance in older Masters athletes. *Age and Ageing* 44: 683-686, 2015.
- 410 3. Brooks SV and Faulkner JA. Skeletal muscle weakness in old age: underlying
411 mechanisms. *Med Sci Sports Exerc* 26: 432-439, 1994.
- 412 4. Brown N and Scurr J. Do women with smaller breasts perform better in long-
413 distance running? *European journal of sport science* 16: 965-971, 2016.
- 414 5. Campbell MJ, McComas AJ, and Petito F. Physiological changes in ageing
415 muscles. *J Neurol Neurosurg Psychiatry* 36: 174-182, 1973.
- 416 6. Carfagno DG and Hendrix JC, 3rd. Overtraining syndrome in the athlete:
417 current clinical practice. *Current sports medicine reports* 13: 45-51, 2014.
- 418 7. Carter SL, Rennie C, and Tarnopolsky MA. Substrate utilization during
419 endurance exercise in men and women after endurance training. *Am J Physiol*
420 *Endocrinol Metab* 280: E898-907, 2001.
- 421 8. Coast JR, Blevins JS, and Wilson BA. Do gender differences in running
422 performance disappear with distance? *Can J Appl Physiol* 29: 139-145, 2004.
- 423 9. Cohen J. Statistical power analysis for the behavioral sciences. Hillsdale, NJ:
424 Lawrence Erlbaum Associates, 1988.
- 425 10. Connick MJ, Beckman EM, and Tweedy SM. Relative age affects marathon
426 performance in male and female athletes. *Journal of Sports Science and*
427 *Medicine* 14: 669-674, 2015.
- 428 11. Coupe C, Svensson RB, Grosset JF, Kovanen V, Nielsen RH, Olsen MR,
429 Larsen JO, Praet SF, Skovgaard D, Hansen M, Aagaard P, Kjaer M, and
430 Magnusson SP. Life-long endurance running is associated with reduced
431 glycation and mechanical stress in connective tissue. *Age (Dordrecht,*
432 *Netherlands)* 36: 9665, 2014.
- 433 12. Crewther BT, Cook C, Cardinale M, Weatherby RP, and Lowe T. Two
434 emerging concepts for elite athletes: the short-term effects of testosterone and
435 cortisol on the neuromuscular system and the dose-response training role of
436 these endogenous hormones. *Sports medicine (Auckland, NZ)* 41: 103-123,
437 2011.
- 438 13. Deaner RO. Distance running as an ideal domain for showing a sex difference
439 in competitiveness. *Archives of Sexual Behavior* 42: 413-428, 2013.
- 440 14. Deaner RO, Addona V, Carter RE, Joyner MJ, and Hunter SK. Fast men slow
441 more than fast women in a 10 kilometer road race. *PeerJ* 2016, 2016.
- 442 15. Deaner RO and Mitchell D. More men run relatively fast in U.S. road races,
443 1981-2006: a stable sex difference in non-elite runners. *Evol Psychol* 9: 600-
444 621, 2011.
- 445 16. Deaner RO and Mitchell D. More men run relatively fast in U.S. road races,
446 1981-2006: a stable sex difference in non-elite runners. *Evolutionary*
447 *psychology : an international journal of evolutionary approaches to*
448 *psychology and behavior* 9: 600-621, 2011.
- 449 17. Drey M, Sieber CC, Degens H, McPhee J, Korhonen MT, Müller K, Ganse B,
450 and Rittweger J. Relation between muscle mass, motor units and type of

- 451 training in master athletes. *Clinical Physiology and Functional Imaging* 36:
 452 70-76, 2016.
- 453 18. Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N,
 454 Powell KE, Bauman A, and Lee IM. Does physical activity attenuate, or even
 455 eliminate, the detrimental association of sitting time with mortality? A
 456 harmonised meta-analysis of data from more than 1 million men and women.
 457 *Lancet*, 2016.
- 458 19. Holmes PS and Wright DJ. Motor cognition and neuroscience in sport
 459 psychology. *Current opinion in psychology* 16: 43-47, 2017.
- 460 20. Hoogkamer W, Kram R, and Arellano CJ. How Biomechanical Improvements
 461 in Running Economy Could Break the 2-hour Marathon Barrier. *Sports*
 462 *medicine (Auckland, NZ)*, 2017.
- 463 21. Hunter SK and Stevens AA. Sex differences in marathon running with
 464 advanced age: physiology or participation? *Med Sci Sports Exerc* 45: 148-156,
 465 2013.
- 466 22. Hunter SK, Stevens AA, Magennis K, Skelton KW, and Fauth M. Is there a
 467 sex difference in the age of elite marathon runners? *Med Sci Sports Exerc* 43:
 468 656-664, 2011.
- 469 23. Jokl P, Sethi PM, and Cooper AJ. Master's performance in the New York City
 470 Marathon 1983-1999. *Br J Sports Med* 38: 408-412, 2004.
- 471 24. Kayser B. Exercise starts and ends in the brain. *European Journal of Applied*
 472 *Physiology* 90: 411-419, 2003.
- 473 25. Keegan R, Harwood C, Spray C, and Lavallee D. A qualitative investigation of
 474 the motivational climate in elite sport. *Psychology of Sport and*
 475 *ExerciseElsevier Ltd* 2013, pp 97-107.
- 476 26. Kenefick RW and Cheuvront SN. Hydration for recreational sport and
 477 physical activity. *Nutr Rev* 70 Suppl 2: S137-142, 2012.
- 478 27. Knechtle B, Nikolaidis PT, König S, Rosemann T, and Rüst CA. Performance
 479 trends in master freestyle swimmers aged 25–89 years at the FINA World
 480 Championships from 1986 to 2014. *Age* 38: 1-8, 2016.
- 481 28. Knechtle B, Nikolaidis PT, Rosemann T, and Rüst CA. Performance trends in
 482 3000 m open-water age group swimmers from 25 to 89 years competing in the
 483 FINA World Championships from 1992 to 2014. *Research in Sports Medicine*
 484 25: 67-77, 2017.
- 485 29. Knechtle B, Nikolaidis PT, Rosemann T, and Rust CA. Performance trends in
 486 age group breaststroke swimmers in the FINA World Championships 1986-
 487 2014. *The Chinese journal of physiology* 59: 247-259, 2016.
- 488 30. Knechtle B, Rust CA, Rosemann T, and Lepers R. Age-related changes in
 489 100-km ultra-marathon running performance. *Age (Dordrecht, Netherlands)*
 490 34: 1033-1045, 2012.
- 491 31. Knechtle B, Valeri F, Nikolaidis P, Zingg M, Rosemann T, and Rüst C. Do
 492 women reduce the gap to men in ultra-marathon running? 2016.
- 493 32. Koral J, Oranchuk DJ, Herrera R, and Millet GY. Six Sessions of Sprint
 494 Interval Training improves running performance in trained athletes. *Journal of*
 495 *strength and conditioning research*, 2017.
- 496 33. Kraemer WJ, Ratamess NA, Flanagan SD, Shurley JP, Todd JS, and Todd TC.
 497 Understanding the Science of Resistance Training: An Evolutionary
 498 Perspective. *Sports medicine (Auckland, NZ)*, 2017.

- 499 34. Kremer J and Moran A. Swifter, higher, stronger: The history of sport
500 psychology. *The psychologist*, 2008, pp 740-742.
- 501 35. Kusy K and Zieliński J. Sprinters versus long-distance runners: How to grow
502 old healthy. *Exercise and Sport Sciences Reviews* 43: 57-64, 2015.
- 503 36. Lee DC, Pate RR, Lavie CJ, Sui X, Church TS, and Blair SN. Leisure-time
504 running reduces all-cause and cardiovascular mortality risk. *Journal of the*
505 *American College of Cardiology* 64: 472-481, 2014.
- 506 37. Lepers R and Cattagni T. Do older athletes reach limits in their performance
507 during marathon running? 2011.
- 508 38. Lepers R, Stapley PJ, and Cattagni T. Centenarian athletes: Examples of
509 ultimate human performance? *Age and Ageing* 45: 729-733, 2016.
- 510 39. Lexell J. Human aging, muscle mass, and fiber type composition. *Journals of*
511 *Gerontology - Series A Biological Sciences and Medical Sciences* 50: 11-16,
512 1995.
- 513 40. Leyk D, Erley O, Ridder D, Leurs M, Ruther T, Wunderlich M, Sievert A,
514 Baum K, and Essfeld D. Age-related changes in marathon and half-marathon
515 performances. *Int J Sports Med* 28: 513-517, 2007.
- 516 41. Lippi G, Banfi G, Favaloro EJ, Rittweger J, and Maffulli N. Updates on
517 improvement of human athletic performance: focus on world records in
518 athletics. *Br Med Bull* 87: 7-15, 2008.
- 519 42. Mohlenkamp S, Lehmann N, Breuckmann F, Brocker-Preuss M, Nassenstein
520 K, Halle M, Budde T, Mann K, Barkhausen J, Heusch G, Jockel KH, and
521 Erbel R. Running: the risk of coronary events : Prevalence and prognostic
522 relevance of coronary atherosclerosis in marathon runners. *Eur Heart J* 29:
523 1903-1910, 2008.
- 524 43. Moon JR. Role of muscle mass on sprint performance: Gender differences? by
525 Jorge Perez-Gomez, German Vicente Rodriguez, Ignacio Ara, Hugo
526 Olmedillas, Javier Chavarren, Juan Jose González-Henriquez, Cecilia Dorado
527 and José A. L. Calbet. *European Journal of Applied Physiology* 103: 375,
528 2008.
- 529 44. Mottaghi M, Atarodi A, and Rohani Z. The Relationship between Coaches'
530 and Athletes' Competitive Anxiety, and their Performance. *Iran J Psychiatry*
531 *Behav Sci* 7: 68-76, 2013.
- 532 45. Naranjo JD, Dziki JL, and Badylak SF. Regenerative Medicine Approaches
533 for Age-Related Muscle Loss and Sarcopenia: A Mini-Review. *Gerontology*
534 63: 580-589, 2017.
- 535 46. Paffenbarger RS, Jr., Hyde RT, Wing AL, and Hsieh CC. Physical activity,
536 all-cause mortality, and longevity of college alumni. *N Engl J Med* 314: 605-
537 613, 1986.
- 538 47. Pollock ML, Foster C, Knapp D, Rod JL, and Schmidt DH. Effect of age and
539 training on aerobic capacity and body composition of master athletes. *Journal*
540 *of Applied Physiology* 62: 725-731, 1987.
- 541 48. Proctor DN and Joyner MJ. Skeletal muscle mass and the reduction of
542 $\dot{V}O_{2max}$ in trained older subjects. *Journal of Applied Physiology* 82: 1411-
543 1415, 1997.
- 544 49. Reaburn P and Dascombe B. Anaerobic performance in masters athletes.
545 *European Review of Aging and Physical Activity* 6: 39-53, 2009.
- 546 50. Rittweger J, Di Prampero PE, Maffulli N, and Narici MV. Sprint and
547 endurance power and ageing: An analysis of master athletic world records.
548 *Proceedings of the Royal Society B: Biological Sciences* 276: 683-689, 2009.

- 549 51. Rossetti ML, Steiner JL, and Gordon BS. Androgen-mediated regulation of
550 skeletal muscle protein balance. *Molecular and cellular endocrinology* 447:
551 35-44, 2017.
- 552 52. Salinero JJ, Soriano ML, Lara B, Gallo-Salazar C, Areces F, Ruiz-Vicente D,
553 Abian-Vicen J, Gonzalez-Millan C, and Del Coso J. Predicting race time in
554 male amateur marathon runners. *J Sports Med Phys Fitness*, 2016.
- 555 53. Senefeld J, Joyner MJ, Stevens A, and Hunter SK. Sex differences in elite
556 swimming with advanced age are less than marathon running. *Scand J Med*
557 *Sci Sports* 26: 17-28, 2016.
- 558 54. Senefeld J, Smith C, and Hunter SK. Sex Differences in Participation,
559 Performance, and Age of Ultramarathon Runners. *International journal of*
560 *sports physiology and performance* 11: 635-642, 2016.
- 561 55. Sparling PB and Cureton KJ. Biological determinants of the sex difference in
562 12-min run performance. *Medicine and Science in Sports and Exercise* 15:
563 218-223, 1983.
- 564 56. Tanaka H and Seals DR. Endurance exercise performance in Masters athletes:
565 age-associated changes and underlying physiological mechanisms. *J Physiol*
566 586: 55-63, 2008.
- 567 57. Tarumi T, Gonzales MM, Fallow B, Nualnim N, Lee J, Pyron M, Tanaka H,
568 and Haley AP. Cerebral/Peripheral Vascular Reactivity and Neurocognition in
569 Middle-Age Athletes. *Med Sci Sports Exerc* 47: 2595-2603, 2015.
- 570 58. Thibault V, Guillaume M, Berthelot G, Helou NE, Schaal K, Quinquis L,
571 Nassif H, Tafflet M, Escolano S, Hermine O, and Toussaint JF. Women and
572 Men in Sport Performance: The Gender Gap has not Evolved since 1983. *J*
573 *Sports Sci Med* 9: 214-223, 2010.
- 574 59. Tipton CM. Sports medicine: a century of progress. *J Nutr* 127: 878s-885s,
575 1997.
- 576 60. Tucker R and Santos-Concejero J. "An Imminent Sub 2-Hours Marathon is
577 Unlikely: historical Trends of the Gender Gap in Running Events".
578 *International journal of sports physiology and performance*: 1-18, 2016.
- 579 61. Unterwiesing CM, Knechtle B, Nikolaidis PT, Rosemann T, and Rust CA.
580 Increased participation and improved performance in age group backstroke
581 master swimmers from 25-29 to 100-104 years at the FINA World Masters
582 Championships from 1986 to 2014. *Springerplus* 5: 645, 2016.
- 583 62. Weiss M, Newman A, Whitmore C, and Weiss S. One hundred and fifty years
584 of sprint and distance running - Past trends and future prospects. *European*
585 *journal of sport science* 16: 393-401, 2016.
- 586 63. Wen CP, Wai JP, Tsai MK, Yang YC, Cheng TY, Lee MC, Chan HT, Tsao
587 CK, Tsai SP, and Wu X. Minimum amount of physical activity for reduced
588 mortality and extended life expectancy: a prospective cohort study. *Lancet*
589 378: 1244-1253, 2011.
- 590 64. Whipp BJ and Ward SA. Will women soon outrun men? *Nature* 355: 25, 1992.
- 591 65. Whyte G. Age, sex and (the) race: gender and geriatrics in the ultra-endurance
592 age. *Extrem Physiol Med* 3: 1, 2014.
- 593 66. Williams T, Walz E, Lane AR, Pebole M, and Hackney AC. The effect of
594 estrogen on muscle damage biomarkers following prolonged aerobic exercise
595 in eumenorrheic women. *Biol Sport* 32: 193-198, 2015.
- 596 67. Wright VJ and Perricelli BC. Age-related rates of decline in performance
597 among elite senior athletes. *Am J Sports Med* 36: 443-450, 2008.

- 598 68. Zingg MA, Karner-Rezek K, Rosemann T, Knechtle B, Lepers R, and Rust
599 CA. Will women outrun men in ultra-marathon road races from 50 km to
600 1,000 km? Springerplus 3: 97, 2014.
- 601 69. Zingg MA, Knechtle B, Rust CA, Rosemann T, and Lepers R. Reduced
602 performance difference between sexes in master mountain and city marathon
603 running. International journal of general medicine 6: 267-275, 2013.
- 604

ACCEPTED

605 **Table 1** Available data for each race distance at the events considered in our study

606

| | | | 100 m | 200 m | 400 m | 800 m | 1500 m | 5000 m | 1000 m | marathon |
|------|-------------|-------|-------|-------|-------|-------|--------|--------|--------|----------|
| 1978 | Viareggio | men | x | x | x | x | x | x | x | x |
| | | women | | | | | x | x | | x |
| 1980 | Helsinki | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | | x |
| 1982 | Strasbourg | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 1984 | Brighton | men | x | x | x | x | x | x | x | |
| | | women | x | x | x | x | x | x | | |
| 1986 | Malmö | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 1990 | Budapest | men | x | x | x | x | x | x | x | |
| | | women | | | | | | | | |
| 1994 | Athen | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 1996 | Malmö | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 1998 | Cesenatico | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2000 | Jyväskylä | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2002 | Potsdam | men | x | x | x | x | x | x | x | |
| | | women | x | x | x | x | x | x | x | |
| 2004 | Aarhus | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2006 | Poznan | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2008 | Ljubljana | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2010 | Nyiregyháza | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2012 | Zittau | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |
| 2014 | Izmir | men | x | x | x | x | x | x | x | x |
| | | women | x | x | x | x | x | x | x | x |

607 **Table 2** Coefficients (C) and standard errors of estimate (SEE) from multi-variate
608 regression models for race speed of participants by sex and race distance.
609

| | C | SEE | p |
|-------------------------------|----------|------------|----------|
| Sex | -3.18 | 0.08 | <0.001 |
| Race distance | -2.06 | 0.02 | <0.001 |
| Interaction sexxrace distance | 0.17 | 0.02 | <0.001 |

610
611

Table 3 Coefficients (C) and standard errors of estimate (SEE) from multi-variate regression models for race speed of participants by sex and age group for each race distance.

| | | C | SEE | p |
|-----------------|---------------------------|----------|------------|----------|
| 100 m | Sex | -4.06 | 0.18 | <0.001 |
| | Age group | -1.19 | 0.02 | <0.001 |
| | Interaction sex×age group | -0.10 | 0.03 | 0.001 |
| 200 m | Sex | -4.63 | 0.17 | <0.001 |
| | Age group | -1.29 | 0.02 | <0.001 |
| | Interaction sex×age group | -0.02 | 0.03 | 0.481 |
| 400 m | Sex | -5.21 | 0.15 | <0.001 |
| | Age group | -1.34 | 0.02 | <0.001 |
| | Interaction sex×age group | 0.14 | 0.03 | <0.001 |
| 800 m | Sex | -4.85 | 0.13 | <0.001 |
| | Age group | -1.24 | 0.01 | <0.001 |
| | Interaction sex×age group | 0.23 | 0.03 | <0.001 |
| 1500 m | Sex | -4.37 | 0.14 | <0.001 |
| | Age group | -1.08 | 0.02 | <0.001 |
| | Interaction sex×age group | 0.20 | 0.03 | <0.001 |
| 5000 m | Sex | -4.16 | 0.11 | <0.001 |
| | Age group | -0.98 | 0.01 | <0.001 |
| | Interaction sex×age group | 0.21 | 0.02 | <0.001 |
| 1000 m | Sex | -4.13 | 0.15 | <0.001 |
| | Age group | -0.91 | 0.02 | <0.001 |
| | Interaction sex×age group | 0.27 | 0.03 | <0.001 |
| Marathon | Sex | -4.51 | 0.18 | <0.001 |
| | Age group | -0.82 | 0.02 | <0.001 |
| | Interaction sex×age group | 0.35 | 0.04 | <0.001 |

Table 4 Coefficients (C) and standard errors of estimate (SEE) from multi-variate regression models for race speed of participants by sex and calendar year for each race distance.

| | | C | SEE | p |
|-----------------|-------------------------------|----------|------------|----------|
| 100 m | Sex | -0.87 | 33.20 | 0.979 |
| | Calendar year | 0 | 0.01 | 0.948 |
| | Interaction sex×calendar year | 0 | 0.02 | 0.944 |
| 200 m | Sex | -58.47 | 35.01 | 0.095 |
| | Calendar year | -0.02 | 0.01 | 0.161 |
| | Interaction sex×calendar year | 0.03 | 0.02 | 0.113 |
| 400 m | Sex | 6.67 | 34.19 | 0.845 |
| | Calendar year | -0.01 | 0.01 | 0.636 |
| | Interaction sex×calendar year | 0 | 0.02 | 0.783 |
| 800 m | Sex | 25.81 | 29.66 | 0.384 |
| | Calendar year | -0.01 | 0.01 | 0.219 |
| | Interaction sex×calendar year | -0.01 | 0.01 | 0.348 |
| 1500 m | Sex | 4.77 | 24.40 | 0.845 |
| | Calendar year | -0.02 | 0.01 | 0.003 |
| | Interaction sex×calendar year | 0 | 0.01 | 0.779 |
| 5000 m | Sex | -13.03 | 20.69 | 0.529 |
| | Calendar year | -0.02 | 0.01 | 0.010 |
| | Interaction sex×calendar year | 0.01 | 0.01 | 0.596 |
| 1000 m | Sex | 39.86 | 35.34 | 0.260 |
| | Calendar year | -0.02 | 0.01 | 0.017 |
| | Interaction sex×calendar year | -0.02 | 0.02 | 0.239 |
| Marathon | Sex | -3.69 | 22.42 | 0.869 |
| | Calendar year | -0.02 | 0.01 | 0.002 |
| | Interaction sex×calendar year | 0 | 0.01 | 0.946 |

624 **List of figures**

625

626 **Figure 1** Speed by race distance and sex. Women are depicted by ▲ and men by
627 ●.

628

629 **Figure 2** Relationship between speed and race duration for women and men.
630 Women are depicted by ▲ and men by ●.

631

632 **Figure 3** Speed by age group and sex for each race distance. Women are
633 depicted by ▲ and men by ●.


634

635 **Figure 4** Speed by calendar year and sex for each race distance. Women are
636 depicted by ▲ and men by ●.

637


638 **Figure 5** Sex difference in speed by race distance and calendar year.

639

640 **Figure 1**


641

642

643 **Figure 2**


644

645


646 **Figure 3**

647

648 **Figure 4**



651 Figure 5



652